
The Cornell Neurosurgery "Gang"



ROENTGENOGRAPHY IN THE LOCALIZATION OF BRAIN TUMOR, BASED UPON A SERIES OF ONE HUNDRED CONSECUTIVE CASES.

By GEORGE J. HEUER AND WALTER E. DANDY. (From the Surgical Service of The Johns Hopkins Hospital.)

Roentgenography was first applied to the study of intra-[311] cranial tumors in 1897 by Obici and Bollici¹ and, almost simultaneously, by Oppenheim.² They attempted to show that brain


[From THE JOHNS HOPKINS HOSPITAL BULLETIN, Vol. XXVII, No. 306, August, 1916.]

Heuer worked closely with Walter Dandy.

A REPORT OF SEVENTY CASES OF BRAIN TUMOR.*

By GEORGE J. HEUER and WALTER E. DANDY. (From the Surgical Service of The Johns Hopkins Hospital.)

It is but nonagant to me !!



George Hener

Chief of Surgery Cincinnati 1921-1932

Chief of Surgery New York Hospital 1982-1948

Eugene Pool

t uncle of J. Lawrence Pool) Chief of Private Attendings New York Hospital 1932

Some of Heuer's operative reports still exist.

October 4, 1932

Exploration for Cerebellar Tusor. Finding of Large Noduler Tumor Presenting Above Mein Sten. Redicel Extirpation. Hemostanis.

Dr. Houer

Dr. Neagher

Local and Avertin

Local infiltration of the scalp after the usual shaving and clounsing. Neck prepared with indine and alcohol. Cerebeller position was adequate and exposure good throughout. Usual cross-bow incision from mestoid tip to encircle the skull 3 cm. above the external compital protuberance l cunarh curnaan and

"When I review my own professional life and its many satisfactions, the greatest is not the surgical operations I have performed nor the thousands of patients I have cured but the successful young men whose training I have directed."

George Heuer

